

Vertex-deleted subgraphs and regular factors from regular graph^{*}

Hongliang Lu ^{a†}, Bing Bai ^b, Wei Wang ^a

^aDepartment of Mathematics

Xi'an Jiaotong University, Xi'an 710049, PR China

^bCenter for Combinatorics, LPMC

Nankai University, Tianjin, China

Abstract

Let k , m and r be three integers such that $2 \leq k \leq m \leq r$. Let G be a $2r$ -regular, $2m$ -edge-connected graph of odd order. We obtain some sufficient conditions, which guarantee $G - v$ contains a k -factor for all $v \in V(G)$.

1 Introduction

All graphs considered are multigraphs (with loops) and finite. Let $G = (V, E)$ be a graph with vertex set $V(G)$ and edge set $E(G)$. The number of vertices of a graph G is called the *order* of G and is denoted by n . On the other hand, the number of edges of G is called the *size* of G and is denoted by e . We denote the degree of vertex v in G by $d_G(v)$. For two subsets $S, T \subseteq V(G)$, let $e_G(S, T)$ denote the number of edges of G joining S to T .

Let $c(G)$ and $c_o(G)$ denote the number of components and the number of odd components of G , respectively. Let k be a positive integer. A *k-factor* of a graph G is a spanning subgraph H of G such that $d_H(x) = k$ for every $x \in V(G)$.

Petersen obtained the following theorem, which is chronologically the first result on k -factors in regular graphs.

Theorem 1.1 (Petersen, [5]) *Every 3-regular, 2-connected graph has a 1-factor.*

For 1-factor in arbitrary graphs, we have the following characterization by Tutte [6].

^{*}This work is supported by the Fundamental Research Funds for the Central Universities.

[†]Corresponding email: luhongliang215@sina.com (H. Lu)

Theorem 1.2 (Tutte, [6]) *A graph G has a 1-factor if and only if $c_o(G - S) \leq |S|$ for all $S \subseteq V(G)$.*

The following theorem is well-known Tutte's k -factor Theorem.

Theorem 1.3 (Tutte, [7]) *Let G be a graph and k be an integer. Then G has a k -factor if and only if, for all $D, S \subseteq V(G)$ with $D \cap S = \emptyset$,*

$$\delta_G(D, S) = k|D| + \sum_{x \in S} d_G(x) - k|S| - e_G(D, S) - q_G(D, S; k) \geq 0,$$

where $q_G(D, S; k)$ is the number of components C of $G - (D \cup S)$ such that $e_G(V(C), S) + k|C| \equiv 1 \pmod{2}$. Moreover, $\delta_G(D, S) \equiv k|V(G)| \pmod{2}$. (Sometimes C is called k -odd component.)

The following theorem examines the existence of a 1-factor in vertex-deleted subgraphs of a regular graph.

Theorem 1.4 (Little et al., [2]) *Let G be a $2r$ -regular, $2r$ -edge-connected graph of odd order and u be any vertex of G . Then the graph $G - u$ has a 1-factor.*

Katerinis presented the following result, which generalizes Theorem 1.4.

Theorem 1.5 (Katerinis, [4]) *Let G be a $2r$ -regular, $2r$ -edge-connected graph of odd order and m be an integer such that $1 \leq m \leq r$. Then for every $u \in V(G)$, the graph $G - u$ has an m -factor.*

2 Main result

The purpose of this paper is to present the following result which generalizes Theorem 1.5.

Theorem 2.1 *Let m and r be two integers such that $2 \leq m < r$. Let G be $2r$ -regular, $2m$ -edge-connected graph with odd order. If one of the following conditions holds, then $G - v$ has a k -factor for all $v \in V(G)$.*

- (i) k is even and $2 \leq k \leq m$;
- (ii) k is odd, $3 \leq k \leq m$ and $2m > r$.

Proof. Suppose that the result doesn't hold. Then there exists $u \in V(G)$ such that $G - u$ contains no k -factors. Let $H = G - u$. By Theorem 1.3, there exist two disjoint subsets D and S of $V(G) - u$ such that

$$q_H(D, S; k) + \sum_{x \in S} (k - d_{H-D}(x)) \geq k|D| + 2. \quad (1)$$

Define $S' = S \cup \{u\}$ and $W = (G - D) - S'$.

Claim 1. $c(W) \geq 2$.

Otherwise, suppose $c(W) \leq 1$. We consider two cases.

Case 1. $c(W) = 0$.

Since $c(W) \geq q_H(D, S; k)$, (1) implies

$$\sum_{x \in S} (k - d_{H-D}(x)) \geq k|D| + 2. \quad (2)$$

So $k|S| \geq k|D| + 2$ and hence $|S| > |D|$. But $V(H) = D \cup S$ and $|V(H)|$ is even, therefore

$$|S| \geq |D| + 2. \quad (3)$$

Now since G is $2r$ -regular by Tutte's theorem we have

$$\sum_{x \in S'} (2r - d_{G-D}(x)) \leq 2r|D|,$$

which implies

$$2r|S'| - \sum_{x \in S'} d_{G-D}(x) \leq 2r|D|.$$

Therefore,

$$2r(|S| + 1) - \sum_{x \in S'} d_{G-D}(x) \leq 2r|D|,$$

and hence,

$$(2r - k)|S| + k|S| + 2r - \sum_{x \in S'} d_{G-D}(x) \leq k|D| + (2r - k)|D|. \quad (4)$$

However

$$\begin{aligned} \sum_{x \in S'} d_{G-D}(x) &= \sum_{x \in S} d_{G-D}(x) + d_{G-D}(u) \\ &= \sum_{x \in S} d_{H-D}(x) + e_G(u, S) + d_{G-D}(u). \end{aligned}$$

Therefore (4) becomes

$$k|D| + (2r - k)|D| \geq (2r - k)|S| + k|S| + 2r - \sum_{x \in S} d_{H-D}(x) - e_G(u, S) - d_{G-D}(u) \quad (5)$$

Now using (2) and since $e_G(u, S) \leq d_{G-D}(u) \leq 2r$, (5) implies

$$(2r - k)(|S| - |D|) \leq 2r - 2. \quad (6)$$

Moreover, by (3), $|S| \geq |D| + 2$, we can conclude from (6) that $k \geq r + 1$. That is a contradiction, thus Case 1 can't occur.

Case 2. $c(W) = 1$.

Then we have $q_H(D, S; k) \leq 1$ and this implies

$$\sum_{x \in S} (k - d_{H-D}(x)) \geq k|D| + 1. \quad (7)$$

So $k|S| \geq k|D| + 1$ and hence $|S| > |D|$.

Since G is a $2m$ -edge-connected, $2r$ -regular graph, so we have

$$\begin{aligned} 2r|D| &\geq e_G(D, V(G - D)) = e_G(D, V(W)) + e_G(D, S') \\ &= e_G(D \cup S', V(W)) - e_G(S', V(W)) + e_G(D, S') \\ &= e_G(D \cup S', V(W)) - \left(\sum_{x \in S'} d_{G-D}(x) - 2e_G(S', S') \right) + (2r|S'| - \sum_{x \in S'} d_{G-D}(x)) \\ &\geq e_G(D \cup S', V(W)) - 2 \sum_{x \in S'} d_{G-D}(x) + 2e_G(S', S') + 2r|S'| \\ &\geq 2m - 2 \left(\sum_{x \in S} d_{H-D}(x) + e_G(u, S) + d_{G-D}(u) \right) + 2e_G(S', S') + 2r|S'| \\ &\geq 2m - 2 \sum_{x \in S} d_{H-D}(x) - 2d_{G-D}(u) + 2e_G(S, S) + 2r|S'| \\ &\geq 2m - 2 \sum_{x \in S} d_{H-D}(x) + 2r|S| - 2r. \end{aligned}$$

Now using (7) implies

$$\begin{aligned} 2r|D| &\geq 2m - 2 \sum_{x \in S} d_{H-D}(x) - 2r + 2r|S| \\ &\geq 2m - 2(k|S| - k|D| - 1) - 2r + 2r|S|. \end{aligned}$$

Thus

$$(2r - 2k)(|D| - |S|) \geq 2m - 2r + 2 \geq 2k - 2r + 2$$

from which it follows $|D| \geq |S|$, a contradiction. So Case 2 can't also occur.

So we have $c(W) \geq 2$. We denote the components of W by $C_1, \dots, C_{c(W)}$. Suppose that $e_G(C_1, D \cup S') \leq \dots \leq e_G(C_{c(W)}, D \cup S')$.

Firstly, we consider (i). Then we have

$$\begin{aligned} 2r|D| &\geq e_G(D \cup S', V(W)) - 2 \sum_{x \in S'} d_{G-D}(x) + 2e_G(S', S') + 2r|S'| \\ &\geq (2m - 2)c(W) - 4r + (2r - 2k)|S| + 2r + 2k|D| + 4 \\ &\geq 4m - 2r + (2r - 2k)|S| + 2k|D|. \end{aligned}$$

Thus we have $(2r - 2k)(|D| - |S| + 1) \geq 2m$ from which it follows $|D| \geq |S|$. For every odd component C of W , the integer $k|V(C)| + e_H(V(C), S)$ is odd and since k is an even integer, so $e_H(V(C), S)$ must be odd. Thus $e_H(V(C), S) \geq 1$ and so $\sum_{x \in S} d_{H-D}(x) \geq q_H(D, S; k)$. Hence (1) implies $k|S| \geq k|D| + 2$ from which it follows that $|S| \geq |D| + 1$, a contradiction.

Secondly, we show that (ii), that is, $k \geq 3$ is odd and $2m > r$. By Theorem 1.5, we can assume that $m < r$. So we have

$$\begin{aligned} 2r|D| &\geq e_G(D \cup S', V(W)) - 2 \sum_{x \in S'} d_{G-D}(x) + 2e_G(S', S') + 2r|S'| \\ &\geq (2m - 2)c(W) - 4r + (2r - 2k)|S| + 2r + 2k|D| + 4 \\ &\geq 4m - 2r + (2r - 2k)|S| + 2k|D| \\ &\geq (2r - 2k)|S| + 2k|D| + 2. \end{aligned}$$

Thus we have $(2r - 2k)(|D| - |S|) \geq 4m - 2r \geq 2$ and hence $|D| > |S|$. Let $q = q_H(D, S; k)$. Note that

$$2r|D| \geq 2mq + 2r|S| - 2r - 2 \sum_{x \in S} d_{H-D}(x).$$

So we obtain

$$|D| - |S| \geq \frac{m}{r}q - 1 - \frac{1}{r} \sum_{x \in S} d_{H-D}(x). \quad (8)$$

By (1), we have

$$|D| - |S| \leq \frac{1}{k}(q - \sum_{x \in S} d_{H-D}(x) - 2), \quad (9)$$

and $q \geq k + 2$ since $|D| > |S|$. By (8) and (9), we have

$$\begin{aligned}
0 \leq \left(\frac{1}{k} - \frac{1}{r}\right) \sum_{x \in S} d_{H-D}(x) &\leq \frac{q}{k} - \frac{2}{k} - \frac{mq}{r} + 1 \\
&< \frac{q}{k} - \frac{2}{k} - \frac{q}{2} + 1 \\
&\leq q\left(\frac{1}{k} - \frac{1}{2}\right) - \frac{2}{k} + 1 \\
&\leq (k+2)\left(\frac{1}{k} - \frac{1}{2}\right) - \frac{2}{k} + 1 \\
&= 1 - k/2 < 0,
\end{aligned}$$

a contradiction. We complete the proof. \square

The bounds are sharp. Firstly, we show that the upper bound is sharp. Let G_1 be the complete graph K_{2r+1} from which a matching of size m is deleted. Let G_2 be the bipartite graph with bipartition (U, W) obtained by deleting a matching of size m from $K_{2r, 2r}$. Let G be the $2r$ -regular graph obtained by matching the $2m$ vertices of degree $2r - 1$ in G_1 to $2m$ vertices of degree $2r - 1$ in G_2 . Clearly, G is $2m$ -edge-connected. Let $m^* \geq m + 1$. Now we show that $G - u$ contains no m^* -factors for all $u \in U \cup W$. Without loss generality, suppose that $u \in U$. Let $D = U - u$, $S = W$ and $G' = G - u$. Note that $q_{G'}(D, S; m^*) = 1$ if $m^* \not\equiv m \pmod{2}$ and $q_{G'}(D, S; m^*) = 0$ if $m^* \equiv m \pmod{2}$. Then we have

$$m^*|D| - m^*|S| + \sum_{x \in S} d_{G'-D}(x) - q_{G'}(D, S; m^*) \leq -2 < 0.$$

So by Theorem 1.3, $G - u$ contains no m^* -factors.

Next we show that the lower bound is sharp. Let Γ be the complete graph K_{2r+1} from which a matching of size $r - 1$ is deleted. Take $2r - 1$ disjoint copies of Γ . Let M_{r-1} be a matching of size $r - 1$. Let H be the $2r$ -regular graph obtained by matching the $2r - 2$ vertices of degree $2r - 1$ in each of $2r - 1$ disjoint copies of Γ to the vertex set S of M_{r-1} . Since $c_o(H - V(M_{r-1})) = 2r - 1 > |V(M_{r-1}) - v| = 2r - 3$ for all $v \in V(M_{r-1})$, so by Theorem 1.2, $G - v$ contains no 1-factor for all $v \in V(M_{r-1})$. So the lower bound is sharp.

Finally, we show that the condition $2m > r$ is sharp. Otherwise, suppose that $2m \leq r$. Let R_1 denote the complete bipartite graph $K_{2r, 2r-1}$ with bipartition (U, W) , where $|U| = 2r$ and $|W| = 2r - 1$. Let R_2 be the complete graph K_{2r+1} from which a matching of size m is deleted. Take two copies of R_2 . Match $4m$ vertices of degree $2r - 1$ of two copies of R_2 to $4m$ vertices of degree $2r - 1$ of $K_{2r, 2r-1}$, and then add a matching of size $r - 2m$ to the rest vertices of degree $2r - 1$ of $K_{2r, 2r-1}$. Then we obtain a $2r$ -regular, $2m$ -edge-connected graph R . Let $u \in U$ and $R' = R - u$. Let $D = U - u$ and $S = W$. Since k is odd, so

$q_{R'}(D, S; k) = 2$ and hence we have

$$k|D| - k|S| + \sum_{x \in S} d_{R'-D}(x) - q_{G'}(D, S; k) = -2 < 0.$$

So by Theorem 1.3, $G - u$ contains no k -factors.

References

- [1] L. Collatz and U. Sinogowitz, Spektren endlicher Grafen, *Abh. Math. Sem. Univ. Hamburg*, **21** (1957), 63-77.
- [2] C. H. C. Little, D. D. Grant and D. A. Holton, On defect-d matchings in graphs, *Discrete Math.*, **13** (1975), 41-54.
- [3] L. Lovász, Subgraphs with prescribed valencies, *J. Combin. Theory*, **8** (1970), 391-416.
- [4] P. Katerinis, Regular factors in vertex-deleted subgraphs of regular graphs, *Discrete Math.*, **131** (1994), 357-361.
- [5] J. Petersen, Die Theorie der regulären Graph, *Acta Math.*, **15** (1891), 314-328.
- [6] W. T. Tutte, The factorizations of linear graphs, *J. London Math. Soc.*, **22** (1947), 107-111.
- [7] W. T. Tutte, The factors of graphs, *Canad. J. Math.*, **4** (1952), 314-328.